

# Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

1911 October 30, 1908

first ten days of the current month is 62, with 15 deaths. I believe, however, that conditions are better than these figures would indicate. So far as my opportunities for observation permit me to judge, I believe that at least 50 per cent of these reported cases were estivo-autumnal fever. Of the latter disease very few were fatal, so that the number of deaths (15) reported as due to yellow fever is fairly near the truth and so raises the yellow-fever mortality rate to about 50 per cent, which I believe to be fairly accurate, and if anything rather under than over states it.

#### NICARAGUA.

### Report from Bluefields, fruit port.

Acting Assistant Surgeon Goodman reports, October 11, 1903, as follows: Week ended October 10, 1903. Present officially estimated population, 4,000; number of deaths during the week, 3; prevailing disease, malarial fever; general sanitary condition of this port and the surrounding country during the week, good. Two of the above-reported deaths were from gunshot wounds, the third from tuberculosis.

Bills of health were issued to the following-named vessels:

Date.	Name of vessel.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of baggage disinfected.
Oct. 4	Alabama	18	5	0	0
8	Utstein	15	1	0	2
8	Agnes	13	0	0	0
9	Osceola	32	0	0	0

### QUEENSLAND.

## Plague in Brisbane and Townsville.

## Consul-General Bray transmits the following telegrams:

VICTORIA, September 15, 1903.

Premier, Queensland, reports two cases plague occurred Brisbane, which ended fatally 12th instant. All precautions taken.

EDMUND BARTON.

Victoria, September 10, 1903.

Premier, Queensland, reports fresh case plague occurred Townsville 4th instant; ended fatally 6th. All precautions taken.

EDMUND BARTON.

Foreign and insular statistical reports of countries and cities—Yearly and monthly.

Australia—New South Wales—Newcastle.—Month of August, 1903. Estimated population, 49,100. Total number of deaths, 50, including enteric fever 1, and 4 from scarlet fever.

Queensland—Brisbane.—Month of May, 1903. Estimated population, 119,428. Total number of deaths, 137, including diphtheria 2, enteric fever 4, and 4 from plague.